

Molecular Imaging – Emerging Techniques and Staging of Prostate Cancer

Michael Pacold, HMS III

Gillian Lieberman, MD

Non Nova. . .Sed Novae (Not New Things, but Things Done Newly)

- Molecular Imaging: defined as visualization of a unique process *in vivo*, using a specific probe and imaging modality
- Not a new field: same principle as Nuclear Medicine, but more specific
- May be viewed as a “Special Stain” for the Radiologist


Research application: Imaging a Plaque in the Brain

- Alzheimer's disease: 4 million cases in US
- At present, no drugs to prevent or disrupt plaque formation
- Disease evaluation requires neuropathology
- *In vivo* plaque detection useful for drug development (would allow rapid detection of plaques in experimental animals)


Michael Pacold, HMS III

Gillian Lieberman, MD

An *In vivo* Congo Red



Congo Red



AOI-987

Modality: Optical
(Near-Infrared
Fluorescence)

Michael Pacold, HMS III

Gillian Lieberman, MD


Hintersteiner *et al.*, *Nat. Biotech* **2005**, 23, 577-583

Alzheimer's plaques in a mouse

APP23 + AOI-987
(Fluorescence visible at
240 min.)

Wild Type + AOI-987
(no fluorescence at
240 min.)

APP23 + Saline
(no fluorescence)



Michael Pacold, HMS III



Gillian Lieberman, MD

Hintersteiner *et al.*, *Nat. Biotech* **2005**, 23, 577-583

Translational application: Imaging a Plaque in the Aorta


- Coronary Artery Disease: ~500,000 deaths/year
- Conventional imaging assesses calcification and degree of stenosis – but not risk of plaque rupture
- Vulnerable plaque cannot be imaged at present – but has high levels of inflammatory cell and protease activity that might be imaged

An active probe for Cathepsin B




Cathepsin B activity – *in vivo*

MRI



Fluorescence-mediated tomography-shows active
Cathepsin B



Michael Pacold, HMS III


Gillian Lieberman, MD

Chen *et al.*, *Circulation* **2002**, 105, 2766-2771


Lymph Node Imaging for Prostate Cancer

- ~200,000 diagnoses/year; ~30,000 deaths
- Two principal avenues of therapy based on metastases
 - Local: Radical prostatectomy, radiotherapy, or watchful waiting
 - Locally advanced/metastatic: Androgen-deprivation therapy and radiation
- Current methods to detect metastases limited by node size

Superparamagnetic Iron Nanoparticles – a marker for the lymph system



Electron Micrograph



Model of Nanoparticle coated with 10 kDa dextran


Modality: MRI (lymphatics have low signal 24 hrs. post injection with nanoparticles)

Michael Pacold, HMS III

Gillian Lieberman, MD

Harisinghani *et al.*, *NEJM* **2003**, 348, 2491-2499¹⁰

Imaging the Lymphatic System




Michael Pacold, HMS III


Gillian Lieberman, MD

Harisinghani *et al.*, *NEJM* 2003, 348, 2491-2499 ¹¹

Metastases Visualized *in vivo*



Pathology post-excision



Michael Pacold, HMS III

Gillian Lieberman, MD

Harisinghani *et al.*, *NEJM* **2003**, 348, 2491-2499¹²

Lymph Node Metastasis Detection

All sizes	MRI	MRI+Probe
Sensitivity	35.4	90.5*
Specificity	90.4	97.8
5-10 mm		
Sensitivity	28.5	96.4*
Specificity	87.2	99.3
<5 mm		
Sensitivity	0	41.1
Specificity	100	98.1

Michael Pacold, HMS III

Gillian Lieberman, MD

*P<0.001

Harisinghani *et al.*, *NEJM* 2003, 348, 2491-2499¹³

Conclusions

- Molecular imaging of specific biological processes enhances the diagnostic power of radiology
- At a basic research level, molecular imaging is useful for observing molecular events in living organisms and for evaluation of therapeutics
- The clinical applications of molecular imaging include earlier detection and more effective intervention and treatment

References

- Chen, J., *et al.* *In vivo* imaging of proteolytic activity in atherosclerosis. *Circulation* **2002**, *105*, 2766-2771.
- Gross, S. and Piwnica-Worms, D. Spying on cancer: Molecular imaging *in vivo* with genetically encoded reporters. *Cancer Cell* **2005**, *7*, 5-15.
- Harisinghani, M., *et al.* Noninvasive detection of clinically occult lymph-node metastases in prostate cancer. *N. Engl. J. Med.* **2003**, *348*, 2491-2499.
- Harisinghani, M. and Weissleder, R. Sensitive, noninvasive detection of lymph node metastases. *PLoS Med.* **2004**, *1*, e66.
- Hintersteiner, M. *et al.* *In vivo* detection of amyloid- β deposits by near-infrared imaging using an oxazine-derivative probe. *Nat. Biotech.* **2005**, *23*, 577-583.
- Jaffer, F.A. and Weissleder, R. Molecular imaging in the clinical arena. *JAMA* **2005**, *293*, 855-862.
- Mahmood, U., *et al.* Near-Infrared Optical Imaging of Protease Activity for Tumor Detection. *Radiology* **1999**, *213*, 866-870.
- Ntziachristos, V., *et al.* Looking and listening to light: the evolution of whole-body photonic imaging. *Nat. Biotech.* **2005**, *23*, 313-320.

Michael Pacold, HMS III

Gillian Lieberman, MD

Acknowledgements

- Gillian Lieberman, MD
- Larry Barbaras
- Pamela Lepkowski

Michael Pacold, HMS III

Gillian Lieberman, MD